

Lisbon Falls North Quadrangle, Maine

Surficial geologic mapping by

Thomas K. Weddle
Amanda E. Normand
Alexa A. Bernotavicz

Digital cartography by:
Robert A. Johnston

Robert G. Marvinney
State Geologist

Cartographic design and editing by:
Robert D. Tucker

Funding for the preparation of this map was provided in part by the U.S. Geological Survey STATEMAP Program, Cooperative Agreement No. 98HQAG2052.



Maine Geological Survey


Address: 22 State House Station, Augusta, Maine 04333
Telephone: 207-287-2801 E-mail: mgs@maine.gov
Home page: http://www.maine.gov/doc/nr/mc/nr/mc.htm

Open-File No. 03-14

2003

This map supersedes
Open-File Map 99-9.

Surficial Geology




SOURCES OF INFORMATION

Surficial geologic mapping by Thomas K. Weddle and Amanda E. Normand completed during the 1998 field season; funding for this work provided by the U.S. Geological Survey STATEMAP program. Surficial geologic mapping for portions of the quadrangle by Alexa A. Bernotavicz completed during 1993 field season as part of a senior thesis, Bates College, Lewiston, Maine.



Quadrangle Location

SCALE 1 : 24,000



CONTOUR INTERVAL 10 FEET



Topographic base from U.S. Geological Survey Lisbon Falls North quadrangle, scale 1:24,000 using standard U.S. Geological Survey topographic map symbols.

The use of industry, firm, or local government names on this map is for location purposes only and does not implicate responsibility for any present or potential effects on the natural resources.

- af** Artificial fill - Includes landfills, highway and railroad embankments, and dredge spoil areas. These units are mapped only where they are wettable using the contour lines on the map, or where they define the limits of resolvable units. Minor artificial fill is present in virtually all developed areas of the quadrangle.
- Ha** Stream alluvium - Gray to brown fine sand and silt with some gravel. Comprises flood plains along present streams and rivers. Extent of alluvium approximates areas of potential flooding.
- Hw** Freshwater wetlands - Muck, peat, silt, and sand. Poorly drained areas, often with standing water.
- Pa** Braided-stream alluvium - Pleistocene alluvium consisting of fluviially deposited sand and gravel; trough-crossbeds with rare mud drapes and intracasts are found in exposures or excavations in this unit.
- Pmn** Marine nearshore deposits - Pleistocene gravel, sand, and mud deposited as a result of wave activity in nearshore or shallow-marine environments; not associated with beach morphology.
- Pp** Presumpscot Formation - Massive to laminated silty clay with rare dropstones and occasional shelly horizons, which overlie rock and till, and are interbedded with and overlie end moraines and marine fan deposits; includes sand deposited as a distal unit of submarine fans.
- Pem** End moraines - Linear ridges consisting of bedded sand and gravel interbedded with Presumpscot Formation silty clay and overlain by till on the ice-proximal faces of the moraines. Some moraines, or groups of moraines, have been assigned a unique geographic name listed below:
 - Pemsr - Sabattus River moraines
 - Pemwc - Webster Corner moraines
 - Pemr - Ridge Road moraine
 - Pemfb - Fisher Brook moraines
 - Pemlf - Little Falls moraines
 - Pemh - Hudson Road moraines
- Pmd** Marine delta - Glacial-marine delta composed primarily of sorted and stratified sand and gravel. Deposit was graded to surface of late-glacial sea and is distinguished by flat top and foreset and topset beds. Deltas have been assigned a unique geographic name listed below:
 - Pmdph - Pleasant Hill delta; topset-foreset contact at elevation 323 feet (Thompson and others, 1989).
 - Pmdpb - Purington Brook delta; topset-foreset contact at elevation 298 feet (Kettlebottom Road delta of Thompson and others, 1989).

- Pmf** Submarine outwash fans - Thick sand and gravel accumulations formed at the mouth of subglacial tunnels along the receding late Pleistocene ice margin. The sand and gravel is interbedded with and overlain by Presumpscot Formation clay at the distal edges of the fans, and interlayered with and overlain by till at their ice-contact faces. Some fans, or groups of fans have been assigned a unique geographic name listed below:
 - Pmfsp - Sabattus Pond fans
 - Pmfmb - Maxwell Brook fans
 - Pmfsr - Sabattus River fans
 - Pmfk - Kettle bottom fan
 - Pmfwb - West Bowdooin fan
 - Pmfhc - Higgins Corner fan
 - Pmfwh - Whites Hill fan
- Pgi** Ice-contact deposits - Sand and gravel deposited against remnant masses of glacial ice; massive to well stratified; commonly has collapse features and irregular topography.
- Pt** Till - Gravelly to bouldery, sandy, or silty matrix diamiction.
- Ptd** Thin-drift areas - Areas with generally less than ten feet of drift covering bedrock. Till overlies bedrock on hillslopes and ridge crests. Presumpscot Formation silty clay is present in depressions; and nearshore deposits overlie till. Presumpscot Formation, and bedrock on hillslopes and at the base of these slopes. Small rock outcrops, and areas of numerous small outcrops are shown as gray areas.

- Contact between units, dashed where inferred.**
- 135** Glacial striations or grooves - observations made at dot. Number indicates azimuth (in degrees) of ice-flow direction. Where two directions are observed in the same outcrop, flags indicate older trends where discerned.
- End moraine - Ridge of till, sand, and gravel deposited and/or deformed by glacial ice, often mantled by Presumpscot Formation.
- Meltwater channel - Channel eroded by meltwater or later meteoric runoff.
- Ice margin position - Line shows approximate position of ice margin during glacial retreat for major ice-margin positions.
- Drumlin or glacially streamlined hill.
- 135** Chattermarks - Friction cracks that are zones of crescentic fractures on the bedrock surface and are formed by ice or debris in the ice exerting force on the bedrock sufficient to cause it to fracture.

USES OF SURFICIAL GEOLOGY MAPS

A surficial geology map shows all the loose materials such as till (commonly called hardpan), sand and gravel, or clay, which overlie solid ledge (bedrock). Bedrock outcrops and areas of abundant bedrock outcrops are shown on the map, but varieties of the bedrock are not distinguished (refer to bedrock geology map). Most of the surficial materials are deposits formed by glacial and deglacial processes during the last stage of continental glaciation, which began about 25,000 years ago. The remainder of the surficial deposits are the products of postglacial geologic processes, such as river floodplains, or are attributed to human activity, such as fill or local land-modifying features.

The map shows the areal distribution of the different types of glacial features, deposits, and landforms as described in the map explanation. Features such as striations and moraines can be used to reconstruct the movement and position of the glacier and its margin, especially as the ice sheet melted. Other ancient features may include landforms which may record a specific type of environment or climate, now long gone from the state. This glacial geologic history of the quadrangle is useful to the larger understanding of past earth climate, and how our region of the world underwent recent geologically significant climatic and environmental changes. We may then be able to use this knowledge in anticipation of future similar changes for long-term planning efforts, such as coastal development or waste disposal.

Surficial geology maps are often best used in conjunction with related maps such as surficial materials maps or significant sand and gravel aquifer maps for anyone wanting to know what lies beneath the land surface. For example, these maps may aid in the search for water supplies, or economically important deposits such as sand and gravel for aggregate or clay for bricks or pottery. Environmental issues such as the location of a suitable landfill site or the possible spread of contaminants are directly related to surficial geology. Construction projects such as locating new roads, excavating foundations, or siting new homes may be better planned with a good knowledge of the surficial geology of the site. Refer to the list of related publications below.

OTHER SOURCES OF INFORMATION

1. Weddle, T. K., Normand, A. E., Bernotavicz, A. A., and Neil, C. D., 1999, Surficial materials of the Lisbon Falls North quadrangle, Maine: Maine Geological Survey, Open-File Map 99-10.
2. Neil, C. D., 1999, Significant sand and gravel aquifers of the Lisbon Falls North quadrangle, Maine: Maine Geological Survey, Open-File Map 99-23.
3. Thompson, W. B., 1979, Surficial geology handbook for coastal Maine: Maine Geological Survey, 68 p., (out of print).
4. Thompson, W. B., and Borns, H. W., Jr., 1985, Surficial geologic map of Maine: Maine Geological Survey, scale 1:500,000.
5. Thompson, W. B., Crossen, K. J., Borns, H. W., Jr., and Andersen, B. G., 1989, Glaciomarine deltas of Maine and their relation to late Pleistocene-Holocene crustal movements, in Anderson, W. A., and Borns, H. W., Jr. (eds.), Neotectonics of Maine: Maine Geological Survey, Bulletin 40, p. 43-67.